

Nitrogen Immobilization and Mineralization as Affected by Fertilizer Application in Soils

S.A.M. El-Sayed

*Soil and Water Science Department, Faculty of Agriculture,
Al-Azhar University, Assiut, Egypt.*

MINERALIZATION and immobilization processes were studied in soil samples treated with ammonium sulphate, calcium ammonium nitrate, urea, urea ammonium phosphate and/or nitrophosphate along with unhumified dung. The conversion of fertilizer nitrogen into non-KCL extractable fractions occurred within sixteen days and was relatively rapid in the samples treated with urea and urea ammonium phosphate. The mineral nitrogen content during incubation studies was highest in calcium ammonium nitrate treated soils. Amino acid nitrogen appeared to be the main fraction involved in the immobilization and mineralization of nitrogen in soil. Dung behaved as an efficient nitrification inhibitor and slowed the release of nitrogen from fertilizers.

Keywords: Mineralization, Immobilization, Dung, Nitrification inhibitor, Assimilation, Microorganisms.

The conversion of inorganic nitrogen to organic form through assimilation by microorganisms during the decomposition of organic residues is an important aspect of the nitrogen cycle in soil (Negm *et al.*, 1998; El-Douby *et al.*, 2001 and El-Amry *et al.*, 2001). The quantity of nitrogen required to meet the needs of the microbial population depends on the nature of the organic amendments, kinds of nitrogen supplied to the microorganisms, moisture, temperature, pH and aeration (Broadbent, 1968; Broadbent Nakashima, 1970; Kai *et al.*, 1969 and Overrein, 1970). The present investigation deals with immobilization and release of nitrogen in soil samples treated with nitrogenous fertilizers such as ammonium sulphate, calcium ammonium nitrate, urea, urea ammonium

phosphate and nitrophosphate and unhumified dung. The rates and amounts of change were estimated from chemical characteristics of newly immobilized and subsequently released nitrogen in the various nitrogen fractions in acid hydrolysate of soil samples at different stages of incubation. Also, various workers (Huber *et al.*, 1969 and Locascio & Fiskell, 1971) have tried to increase the efficiency of utilization of nitrogenous fertilizers through devices such as the coating of fertilizers and use of nitrification inhibitors (Hamouda *et al.*, 1999; Badran *et al.*, 2000; Habib *et al.*, 2001 and Farghly, 2001). Incorporation of partially decomposed dung, a cheaper source of nitrogen, along with fertilizers in soil may also have good prospects (Arafat, 1998; Hassouna & Hassancin, 2000; Shalan *et al.*, 2001; Allam *et al.*, 2001; El-Sayed 1995; El-Sayed 1997 a&b and El-Sayed, 1998).

Material and Methods

The study was made on a sandy loam soil (0-15 cm) of the Faculty of Agriculture Farm in Assiut, El-Azhar University, which fits the order of Aridisols (irrigated), the suborder of Orthids and the great group of Calciorthids, according to the 7th Approximation system of soil classification (Soil Survey Staff, 1994). The soil contained 0.08 and 0.50% of nitrogen and carbon, respectively and had a pH of 8.6. Electrical conductivity of the saturation extract was 1.5 dS m⁻¹. The soil sample was passed through a 2-mm sieve and subsamples used for the study.

Incubation and determination

Air-dried samples (50 g) were weighed into 100 ml Erlenmeyer flasks and treated with 2.4 g of unhumified dry dung (containing 0.833% N and 30.4% C to give a C/N ratio 36.9) and 200 ppm of nitrogen in the form of ammonium sulphate, urea, calcium ammonium nitrate, urea ammonium phosphate (28% N and 28% P₂O₅) and nitrophosphate, (20% N and 20% P₂O₅). There were two more treatments, namely, (soil + dung) and (soil + fertilizer). The flasks were loosely covered with polyethylene film and placed in an incubator maintained at 30°C for periods of 16, 32, 50, 70, 80, and 110 days. Moisture content in all treatments was maintained at 60% of maximum water holding capacity (Chapman and Pratt, 1961). Ammonium and nitrate nitrogen produced at the different incubation intervals were extracted with 2 N KCl and determined using MgO and Devardas alloy in a specially designed micro-distillation apparatus (Srivastava, 1969; FAO, 1980 and Cottenie, 1980). The procedure adopted for microbially immobilized nitrogen was a modification of the Van Slyke method of protein analysis as followed by Stewart *et al.* (1963a); Page *et al.* (1982) and

Evenhuis & DeWaard (1978). After the mineralized nitrogen was extracted with 2 N KCl, a sample was hydrolyzed with 6 N HCl. Distillable acid-soluble nitrogen (designated as amino sugar $+NH_4-N$) was determined by distilling the filtrate with excess NaOH. Total filtrate (hydrolyzable) nitrogen was determined by the conventional Kjeldahl procedure. The difference between these two was designated as an amino acid nitrogen (non-distillable acid-soluble N). Data were statistically analysed according to Steel & Torrie (1982) and SAS (1988).

Results and Discussion

The results of nitrogen mineralization (ammoniacal and nitrate nitrogen forms) and immobilization (hydrolysable ammonium, amino-sugar and amino-acid forms of nitrogen) as affected by fertilizer application at different incubation intervals, with and without unhumified dung, are presented in Tables 1-4 inclusive.

TABLE 1. Mineralization of nitrogen as affected by different fertilizer application (ppm).

Soil Treatments	Incubation time (days)								Average
	16	32	50	70	80	100	110		
Control	NH ₄ -N	23.5	19.5	20.1	31.3	28.5	21.8	13.0	22.5
	NO ₃ -N	22.6	26.5	26.1	26.2	44.5	63.4	84.4	42.0
	total	46.1	46.0	46.2	57.5	73.0	85.2	97.4	
Ammonium Sulphate	NH ₄ -N	9.3	5.8	9.9	14.1	18.4	20.1	14.4	13.1
	NO ₃ -N	52.1	73.1	94.0	108.1	125.0	160.5	197.1	115.7
	total	61.4	78.9	103.9	122.2	143.4	180.6	211.5	
Urea	NH ₄ -N	0.8	6.0	13.3	13.6	13.9	10.6	6.5	9.2
	NO ₃ -N	36.5	58.2	100.1	143.6	176.0	197.1	218.5	132.9
	total	37.3	64.2	113.4	157.2	189.9	207.7	225.0	
Calcium Ammonium Nitrate	NH ₄ -N	9.8	16.0	15.1	16.7	16.9	14.6	12.6	14.5
	NO ₃ -N	37.1	71.6	84.2	137.3	172.7	200.1	226.0	132.7
	total	46.9	87.6	99.3	154.0	189.6	214.7	238.6	
Urea Ammonium Phosphate	NH ₄ -N	1.8	7.8	4.0	16.6	7.0	8.1	10.0	7.9
	NO ₃ -N	38.9	59.3	105.5	127.6	158.1	180.1	205.6	125.0
	total	40.7	67.1	109.5	144.2	165.1	188.2	215.6	
Nitrophosphate	NH ₄ -N	17.6	7.9	35.0	9.2	43.9	35.1	2.5	21.6
	NO ₃ -N	42.4	56.2	80.9	151.0	131.0	152.6	198.8	116.1
	total	60.0	64.1	115.9	160.2	174.9	187.7	201.3	
Average	NH ₄ -N	10.5	10.5	16.2	16.9	21.4	18.4	9.8	
	NO ₃ -N	38.3	57.5	81.8	115.6	134.6	159.0	188.4	
	total	48.8	68.0	98.0	132.5	156.0	177.4	198.2	

LSD_{0.05} for NH₄-N = 7.8 NO₃-N 23.0

TABLE 2. Effect of dung on the mineralization of nitrogen (ppm).

Soil Treatments		Incubation time (days)							Average
		16	32	50	70	80	100	110	
Dung	NH ₄ -N	22.8	16.6	16.8	25.1	33.6	50.1	13.6	25.5
	NO ₃ -N	7.5	10.0	11.7	20.5	23.6	27.1	83.4	26.3
	total	30.3	26.6	28.5	45.6	57.2	77.2	97.0	
Dung + Ammonium	NH ₄ -N	13.3	22.9	26.3	25.1	24.9	24.4	18.2	22.2
Sulphate	NO ₃ -N	27.5	21.0	22.9	25.6	27.3	71.5	115.7	44.5
	total	40.8	43.9	49.2	50.7	52.2	95.9	133.9	
Dung + Urea	NH ₄ -N	3.2	8.6	8.6	23.7	26.2	24.6	14.1	15.6
	NO ₃ -N	20.0	31.4	44.0	41.1	50.4	109.6	115.1	58.8
	total	23.2	40.0	52.6	64.8	76.6	134.2	129.2	
Dung + Calcium	NH ₄ -N	7.6	20.1	20.6	23.0	21.1	20.7	7.5	17.2
Ammonium	NO ₃ -N	20.1	29.2	50.5	61.8	69.1	104.0	150.7	69.3
Nitrate	total	27.7	49.3	71.1	84.8	90.2	124.7	158.2	
Dung + Urea	NH ₄ -N	9.0	7.5	12.7	8.8	4.8	11.1	12.1	9.4
Ammonium	NO ₃ -N	17.3	30.0	30.8	45.1	57.6	101.9	137.0	60.0
Phosphate	total	26.3	37.5	43.5	53.9	62.4	113.0	149.1	
Dung+Nitrophosphate	NH ₄ -N	1.5	5.1	9.0	7.3	11.4	10.1	9.0	7.6
	NO ₃ -N	26.1	18.4	30.8	46.2	86.1	113.1	137.7	65.5
	total	27.6	23.5	39.8	53.5	97.5	123.2	146.7	
Average	NH ₄ -N	9.6	13.5	15.7	18.8	20.3	23.5	12.4	
	NO ₃ -N	19.8	23.3	31.8	40.1	52.4	87.9	123.3	
	total	29.4	36.8	47.5	58.9	72.7	111.4	135.7	

LSD_{0.05} for: NH₄-N=8.9 NO₃-N = N.S.

Immobilization of nitrogen occurred when fertilizer was applied to the soil. It became more pronounced in the presence of unhumified organic matter. Broadbent & Nakashima (1970); Stewart *et al.* (1963b); Negm *et al.* (1998) and Badran *et al.* (2000) also found increased amounts of tagged organic nitrogen when they incorporated tagged mineral nitrogen in soils. The accumulation of mineral nitrogen in a soil is a net effect of two processes which occur
Egypt. J. Soil Sci. 43, No. 4 (2003)

TABLE 3. Effect of dung and nitrogen fertilizers on hydrolysable NH_4^+ amino-sugar-N fraction (ppm) of soil.

Soil + Treatment	Incubation period (days)										Average
	16	32	50	70	80	100	110				
Soil*	213.7	233.7	234.1	252.9	267.7	279.4	300.2			254.5	
Ammonium sulphate*	246.9	260.4	274.3	274.1	278.8	296.1	325.8			279.5	
Urea*	253.4	287.2	292.9	295.6	297.3	308.6	322.7			294	
Calcium ammonium nitrate*	244.4	264.2	292.1	293.5	311.3	318.6	368.6			299	
Urea-ammonium phosphate*	241.4	268.4	278.2	282.6	302.2	306.1	312.2			284.4	
Nitrophosphate*	249.6	272.3	249.2	278.0	258.3	378.1	312.7			285.5	
Average*	241.6	264.4	270.1	279.5	285.9	314.5	323.7				
Dung**	257.4	286.6	289.4	288.1	310.6	328.1	375.6			305.1	
Dung + ammonium sulphate**	305.9	317.3	316.9	325.1	347.2	351.8	363.0			332.5	
Dung + urea**	299.0	296.6	339.6	330.5	366.0	381.8	417.1			347.2	
Dung + calcium ammonium nitrate**	319.6	319.1	344.6	325.2	352.1	354.5	388.7			343.4	
Dung + urea ammonium phosphate**	313.2	321.7	328.5	343.4	360.4	358.1	366.1			341.6	
Dung + Nitrophosphate**	317.7	324.1	341.2	345.9	355.8	368.1	390.2			349	
Average**	302.1	310.9	326.7	326.4	348.7	357.1	383.5				
Average of all treatment	271.9	287.7	298.4	303.0	317.3	335.8	353.6				

LSD_{0.05} = 18.8LSD_{0.05} = 18.6

TABLE 4. Effect of dung and nitrogen fertilizers on amino-acid-nitrogen (ppm) fraction of soil.

Soil + Treatment	Incubation period (days)										Average
	16	32	50	70	80	100	110				
Soil*	355.0	323.0	279.0	231.0	187.0	80.0	49.0				214.9
Ammonium sulphate*	386.0	356.0	328.0	282.0	271.0	80.0	37.0				248.6
Urea*	382.0	317.0	278.0	258.0	162.0	80.0	38.0				216.4
Calcium ammonium nitrate*	470.0	353.0	309.0	283.0	239.0	99.0	36.0				255.6
Urea-ammonium phosphate*	610.0	361.0	342.0	260.0	182.0	77.0	45.0				268.1
Nitrophosphate*	393.0	354.0	332.0	292.0	257.0	80.0	52.0				251.4
Average*	432.7	344.0	311.3	267.7	216.3	82.7	42.8				
Dung**	612.0	594.0	569.0	557.0	247.0	102.0	29.0				387.1
Dung + ammonium sulphate**	715.0	676.0	669.0	625.0	510.0	150.0	129.0				496.3
Dung + urea**	744.0	688.0	562.0	513.0	361.0	118.0	60.0				435.1
Dung + calcium ammonium nitrate**	724.0	657.0	628.0	570.0	354.0	108.0	61.0				443.1
Dung + urea ammonium phosphate**	751.0	656.0	640.0	618.0	587.0	124.0	84.0				494.3
Dung + Nitrophosphate**	735.0	708.0	608.0	595.0	524.0	130.0	71.0				481.6
Average**	713.5	663.2	612.7	579.7	430.5	122.0	72.3				
Average of all treatment	573.1	503.6	462.0	423.7	323.4	102.4	57.6				

* LSD_{0.05} = N.S.** LSD_{0.05} = N.S.

simultaneously but oppose each other, *i.e.*, mineralization and immobilization. The content of mineral nitrogen after an incubation period of 110 days in case of the control was 97.4 ppm. It was increased to 201.3, 211.5, 215.6, 225.0 and 238.6 ppm by the application of 200 ppm of fertilizer nitrogen as nitrophosphate, ammonium sulphate, urea ammonium phosphate, urea and calcium ammonium nitrate, respectively. The highest concentration of mineral nitrogen in samples treated with calcium ammonium nitrate may be partly due to the calcium content of the fertilizer which might have maintained favourable conditions for nitrifying bacteria by controlling the decline in soil pH (Millbank, 1959; El-Sayed, 1997 a&b and El-Sayed, 1998) and partly due to a lesser amount of $\text{NH}_4\text{-N}$ present in this fertilizer. There is already considerable information showing that $\text{NO}_3\text{-N}$ is as a good source of nitrogen as ammonium for the heterotrophic microflora of the soil but that in the presence of NH_4 and NO_3 the microorganisms use ammonium preferentially (Jansson *et al.*, 1955; Broadbent & Nakashima, 1970; Metwally & Khamis, 1998 and Hassouna & Hassanein, 2000). The mineral nitrogen decreased rapidly at first and changed slowly thereafter. Broadbent (1968); Stewart *et al.* (1963 a&b); Khamis & Metwally (1998) and El-Zaher and El-Kafoury (1999) noted that under conditions favouring biological immobilization, there is an initial period of very rapid decrease in the level of inorganic nitrogen which reaches a minimum within two weeks or less, followed by a period of net mineralization of nitrogen after the needs of the soil population are met. The data in Table 1 indicate that the incorporation of fertilizer nitrogen into non KCl extractable fractions occurred relatively quickly in the urea and urea ammonium phosphate treatments, more than in treatments with ammoniacal and nitrate forms of fertilizers. The results are consistent with the findings of Overrein (1970); Beheiry *et al.* (1998) and Hassanein & El-Shebiny (2000).

A small amount of nitrogen could be accounted for as exchangeable ammonium, but most of the mineral nitrogen was in nitrate form. Nitrate nitrogen increased further with increased incubation time. Most of the fertilizer nitrogen was transformed into a form resistant to acid hydrolysis and some of it was converted into hydrolysable ammonium + amino sugar and amino acid forms. Nommik (1957); Aziz *et al.* (1998) and Hassanein and Hassouna (2000) have described as a function of organic matter the situation in which well decomposed organic matter (humic substances) and living or fresh organic residues exist as two interacting organic pools in which applied nitrogen repeatedly cycles and gradually reverts to increasingly more stable forms (El-Sayed, 1995 and Kristensen *et al.*, 2000).

When dung was applied to the samples, the accumulation of mineral nitrogen was reduced and was found to be even lower than in the controls. One interesting feature is that the amounts of mineral nitrogen in dung and (dung + fertilizer) treated samples were almost similar up to 80 days of incubation, though in later treatments, 200 ppm fertilizer nitrogen were added. These results indicate that microorganisms derived some nitrogen from the native organic source and that the conversion of the $\text{NH}_4\text{-N}$ to the $\text{NO}_3\text{-N}$ form under these conditions was as rapid as was the conversion of organic nitrogen to NH_4 . The results correspond to the findings of Walker *et al.* (1956); Arafat (1998) and El-Douby *et al.* (2001) who reported an accelerated mineralization of humus in soil after addition of flesh organic substances. When dung was mixed in the samples along with fertilizer, the amino acid content in the initial stages was found to be higher than in samples treated with dung alone. This may be due to more favourable conditions for microbial growth. Microorganisms decomposing the organic matter multiple rapidly, result in, in vigorous synthesis of microbial protoplasm. The protoplasm has a rather high N content (3-12%), in most cases higher than the decomposing substrate. The mineral nitrogen will, therefore, be used for the formation of that protoplasm and, if the C/N ratio of the substrate is wide, all the mineralized nitrogen will be reabsorbed by microorganisms and converted into amino acids as body building material (Stewart *et al.*, 1963 a; Cheng & Kurtz, 1963; Chu & Knowles, 1966; Hamouda *et al.*, 1999; Allam *et al.*, 2001 and Farghly, 2001). The results are in conformity with the observations of Broadbent (1968) who found less ammonium and non hydrolyzable nitrogen and more amino acid nitrogen during initial periods of fertilizer application followed by their transformation into organic substances of increasing molecular weight and low availability (Malpassi *et al.*, 2000).

After 100 days of incubation, the mineral nitrogen content in samples treated with dung plus fertilizer increased over control. This may be due to narrowing of the C/N ratio of the dung as decay proceeds and the energy supply diminishes. The microbial population decreases because of the reduced food supply and ultimately a new equilibrium is reached. Therefore, if a crop is to receive a continual supply of nitrogen during the period of its growth, it is necessary that the mineralization rate exceeds that of immobilization, the usual situation in nature. The average mineral nitrogen which was 29.4 ppm after 16 days of incubation increased to 135.7 ppm (Table 2) after 110 days. The effect of this increase in mineral nitrogen was observed on the amino acid fraction of nitrogen

which decreased sharply. After 16 days of incubation, the average figures for amino acid content were 432.7 ppm in fertilizer and 713.5 ppm in fertilizer + dung treatments, and these decreased to 42.8 and 72.3 ppm, respectively (Table 4) after 110 days. Hydrolyzable (ammonium + the amino sugar nitrogen fraction) Table 3 showed a slight gain. Thus it seems that the amino acid fraction has been the most important in contributing to the release of nitrogen in mineral form. Stewart *et al.* (1963 a&b); Habib *et al.* (2001) and Shalan *et al.* (2001) found that changes in the nondistillable acid soluble nitrogen fractions of soil hydrolysates (largely amino acid N) were closely related to inverse changes in inorganic nitrogen of the soil and underwent fairly rapid turnover in their experiment. The results further correspond to the findings of Porter *et al.* (1964); Moore & Russell (1968) and El-Amry *et al.*, 2001) who observed a marked decline in amino acid content of a soil after crop harvesting. They believed that amino acid nitrogen in soils might be more susceptible to mineralization than other forms of organic soil nitrogen. Cornfield (1957) also noted earlier that amino acid content of soils is affected by cropping, cultivation and fertilizer application.

Conclusions

The accumulation of mineral nitrogen in a soil is a net effect of two processes which occur simultaneously but oppose each other, *i.e.*, mineralization and immobilization.

Immobilization of nitrogen occurred when fertilizer was applied to the soil. It became more pronounced in the presence of unhumified organic matter.

Dung behaved as an efficient nitrification inhibitor and slowed the release of nitrogen from fertilizers.

References

- Allam, A.I., Nassar, A.M. and Besheit, S.Y. (2001) Nitrogen fertilizer requirements of *Stevia rebaudiana bertonii*, under Egyptian conditions. *Egypt. J. Res.*, **79**(3), 1005.
- Arafat, S.M. (1998) Uptake and distribution of cadmium in cucumber plant as affected by nitrogen form. *Egypt. J. Soil Sci.* **38**(1-4), 441.
- Aziz, M.A., Sallam, M.F.A., El- Gendy, A.M. and Abd El-Moniem, M. (1998) Effect of natural soil conditioners and irrigation conditions on some chemical properties of sandy soils of Inshas and Cucumber yield. *Egypt. J. Soil Sci.* **38**(1-4), 377.

- Badran, N.M., Khalil, M.E.A. and El-Emam, M.A.A.** (2000) Availability of N, P and K in sandy and clayey soils as affected by the addition of organic materials. *Egypt. J. Soil Sci.* **40**(1-2), 265.
- Beheiry, G.G., El-Bagouri, I.H. and Soliman, A.A.** (1998) The residual effects of application of organo/silicate conditioners on some properties of desert soils and yield of coriander crop. *Egypt. J. Soil Sci.* **38** (1-4),123.
- Broadbent, F.E.** (1968) Turnover of nitrogen in soil organic matter. *In: Organic Matter and Soil Fertility.* North HoHand Pubi. Co. Amsterdam, pp. 61-88.
- Broadbent, F.E. and Nakashima, T.** (1970) Nitrogen immobilization in flooded soil. *Soil Sci. Soc. Am. Proc.* **34**, 218.
- Champman, H.D. and Pratt, P.F.** (1961) *Methods of Analysis for Soil, Plants and Waters.* River Side University of California. Division of Agriculture Science.
- Cheng, H.H. and Kurtz, L.T.** (1963) Chemical distribution of added N in soil. *Soil Sci. Soc. Am. Proc.* **27**, 312.
- Chu, J.P.H. and Knowles, R.** (1966) Mineralization and immobilization of N in bacterial cells and in certain soil organic fractions. *Soil Sci. Soc. Am. Proc.* **30**,210.
- Cornfield, A.H.** (1957) Effect of 8 years fertilizer treatment on the protein N content of four cropped soils. *J. Sci. Food Agric.* **8**, 509.
- Cottenie, A.** (1980) Soil and plant testing as basis of fertilizer recommendations. *FAO Soils Bulletin 38/2* FAO, Rome, Italy.
- El-Amry, H.G., Mahdy, A.M. and Abdel-Salam, S. M.**(2001) Efficacy of combination between Mg and N fertilization on the chemical composition and technological properties of wheat grains. *Egypt. J. Agric. Res.* **79**(3), 1085.
- El-Douby, K.A., Ali, E.A., Toaima, S.E.A. and Abd El-Aziz** (2001) Effect of nitrogen fertilizer, defoliation and plant density on maize grain yield. *Egypt. J. Agric. Res.*, **79** (3), 965.
- El-Sayed, S.A.M.** (1995) Effect of cropping and N-fertilization on changes in different forms of nitrogen in an Oases soil. *J. Agric. Sci. Mansoura Univ.*, **20**(1),453.

- El-Sayed, S.A.M.** (1997a) Nitrifying and denitrifying bacteria in rice fields of Oases soil. *Proceedings of the 9th Conference of Microbiology, Cairo, March 25-27, 1997*, pp. 1-17.
- El-Sayed, S.A.M.** (1997b) Ability of forage cowpea strains to denitrification in New Valley Governorate. *Proceeding of the 9th Conference of Microbiology, Cairo, March, 25-27, 1997*, pp. 18-35.
- El-Sayed, S.A.M.** (1998) Denitrification by guar rhizobia strains in New Valley Governorate. *Egypt. J. Agric. Res.* **76** (2), 451.
- El-Zaher, H. and El-Kafoury, A.A.** (1999) Combination of farmyard manure and N-fertilizer on water and N-fertilizer use efficiency and productivity of sunflower in calcareous soil. *Alex. Sci. Exch.*, **20**, (2), 111.
- Evenhuis, B. and DeWaard, P.W.** (1978) *Simplified Methods for Foliar Analysis*. Royal Tropical Institute. Amsterdam, Holland.
- FAO** (1980) Soil and plant testing and analysis. *FAO. Soils Bulletin 38/1*. FAO, Rome, Italy.
- Farghly, B.S.** (2001) Effect of the preceding winter crop and nitrogen fertilization on yield and yield components of maize and sunflower. *Egypt. J. Agric. Res.* **79**, (4), 1423.
- Habib, F.M., Negm, M.A. and Hassan, M.M.** (2001) Composting of sugarbeet residues. 3- The effect of application dose to a calcareous soil on plant growth and nutrient uptake. *Egypt. J. Agric. Res.*, **79** (4), 1263.
- Hamouda, A.M.M., Abdel Rasoul, S.M. and El-Desouky, H.I.** (1999) Shale treatment and its short-term effect on sandy soil characteristics and yield of peanut and wheat. *Egypt. J. Soil Sci.* **39** (2), 187.
- Hassanein, M.A. and El-Shebiny, G.M.** (2000) Contribution of bio-and mineral nitrogen fertilization in sugarbeet yield. *Alex. Sci. Exch.* **21** (2), 129.
- Hassanein, M.A. and Hassouna, M.G.** (2000) Effect of bio-and mineral nitrogen fertilization on sugarbeet yield and quality in the new reclaimed areas at Nubaria region. *Alex. Sci. Exch.* **21** (2), 153.

- Hassouna, M.G. and Hassanein, M.A.** (2000) Sugarbeet yield and quality as affected by biological and mineral nitrogen fertilization in the newly reclaimed areas at Nubaria Region. *Alex. Sci. Exch.* **21** (3), 211.
- Huber, D.M., Murray, G.A. and Grane, J.M.** (1969) Incubation of nitrification as a detergent to nitrogen loss. *Soil Sci. Soc. Am. Proc.* **33**, 975.
- Jansson, S.L., Hallam, M.J. and Bartholomew, W.V.** (1955) Preferential utilization of NH_4 over NO_3 by microorganisms in the decomposition of oat straw. *Plant Soil*, **6**, 382.
- Kai, H., Ahmed, Z. and Herada, T.** (1969) Factors affecting immobilization and release of N in soil and chemical characteristics of the N newly formed. *Soil Sci. Plant Nutr.* **15**, 207.
- Khamis, M.A. and Metwally, S.M.** (1998) Utilization of organic materials inoculated with microbial decomposers as a nitrogen source for wheat in presence of *Azotobacter* in sandy soil. *Egypt. J. Soil Sci.* **38** (1-4), 81.
- Kristensen, H.L., McCarty, G.W. and Meisinger, J.J.** (2000) Effect of soil structure disturbance on mineralization of organic soil nitrogen. *Soil Sci. Soc. Am. J.* **64**, 371.
- Locascio, S.J. and Fiskell, G.A.** (1971) Sulfur coated fertilizer in water melon production *Sulfur Inst. J.* **7**, 7.
- Malpassi, R.N., Kaspar, T.C., Parking, T.B., Cambardella, C.A. and Nubel, N.A.** (2000) Oat and rye root decomposition effects on nitrogen mineralization. *Soil Sci. Soc. Am. J.* **64**, 208.
- Metwally, S.M. and Khamis, M.A.** (1998) Comparative effects of organic and inorganic nitrogen sources applied to a sandy soil on availability of N and wheat yield. *Egypt. J. Soil Sci.* **38** (1-4), 35.
- Milbank, J.W.** (1959) The physiology of nitrification in Kenya highland soils. *Plant Soil*, **112**, 293.
- Moore, A.W. and Russell, J.S.** (1968) Relative constancy of soil N fractions with varying total soil N. *Int. Congr. Soil Sci. 9th, Adelaide.* **2**, 557-566.
- Negm, M.A., Kerlous, R.G. and Besada, Y.B.** (1998) Different sources of nitrogen and *Rhizobium* inoculation effect on peas growing on a calcareous soils. *Egypt. J. Soil Sci.* **38** (1-4), 69.

- Nommik, H.** (1957) Fixation and defixation of ammonium in soils. *Acta. Agric. Scand.* **7**, 395.
- Overrein, L.N.** (1970) Mineralization and immobilization of tracer N in soil. *Plant Soil*, **32**, 207.
- Page, A.L., Miller, R.H. and Keeney, D.R.** (1982) *Methods of Soil Analysis*, part 2: *Chemical and Microbiological Properties*. Amer. Soc. Agron., Madison, Wisconsin, USA.
- Porter, L.K., Stewart, B.A. and Hass, H.J.** (1964) Effect of long time cropping on hydrolysable organic N fractions in some great plant soils. *Soil Sci. Soc. Am. Proc.* **28**, 268.
- SAS Institute Inc.** (1988) *SAS user's guide*. Statistical Analysis Institute. Inc., Cary, NC. 1674 P.
- Shalan, M.N., Abd-Ellatif, T.A., Soliman, S.G.I. and El-Gaawwas, E.O.** (2001) Effect of some chemical and biofertilizer treatments on roselle plants (*Hibiscus sabdariffa*, L). *Egypt. J. Agric. Res.*, **79** (2), 587.
- Soil Survey Staff** (1994) "Key to Soil Taxonomy," USDA, Sixth ed. Washington, D.C. 265 P.
- Srivastava, O.P.** (1969) A simple apparatus for the microdetermination of NH_4 and $\text{NO}_3\text{-N}$ through steam distillation. *Proc. Natl. Acad. Sci.* **38**, 54.
- Steel, R.G.D. and Torrie, G.H.** (1982) *Principles and Procedures of Statistics. A Biometrical Approach*. Mc Graw-Hill Book Co., 625.
- Stewart, B.A., Porter, L.K. and Johnson, D.D.** (1963a) Immobilization and mineralization of nitrogen in several organic fractions of soil. *Soil Sci. Soc. Am. Proc.* **27**, 302.
- Stewart, B.A., Johnson, D.D. and Porter, L.K.** (1963b) The availability of fertilizer N immobilized during decomposition of straw. *Soil Sci. Soc. Am. Proc.* **27**, 656.
- Walker, T.W., Adams, A.F.R. and Orchiston, H.D.** (1956) Fate of labelled nitrate and ammonium nitrogen when applied to grass and clover grown separately and together. *Soil Sci.* **81**, 339.

(Received 6/200)

تمثيل ومعدنة النيتروجين المتأثر بإضافة الأسمدة في التربة

سعيد عباس محمد السيد

قسم علوم الأراضى والمياه - كلية الزراعة بأسيوط - جامعة الأزهر - مصر.

تم دراسة عمليات المعدنة وتمثيل العناصر في التربة المضاف إليها كل من كبريتات الأمونيوم، ونترات الكالسيوم الأمونيوم، واليوريا، وفوسفات اليوريا الأمونيوم، والنيتروفوسفات إلى جانب سماد الروث الجاف.

تتحول الأسمدة النيتروجينية إلى عناصرها الأولية بدون إستخلاصها بكلوريد البوتاسيوم ويحدث في مدى ١٦ يوما، ويكون سريعا نسبيا في العينات المعاملة مثل اليوريا، وفوسفات اليوريا الأمونيوم.

تم دراسة عنصر النيتروجين خلال فترة التحضين ووجد أن نسبته مرتفعة في التربة المعاملة بنترات الكالسيوم الأمونيوم.

توجد الأحماض الأمينية النيتروجينية كمكون أساسى كأجزاء رئيسية معقدة خلال عمليتى المعدنة وتمثيل العناصر في نيتروجين التربة.

يتصرف سماد الروث سلوكا فعالا بواسطة مثبت النترة (التأزت)، حيث يتحرر النيتروجين ببطء من خلال الأسمدة.